

Q5: Are we confident that quantum spin microscopy will work?

A5: The key pieces are in-place: QMOR, HWIL, and experiment

For details: www.mrfm.org

UW Quantum System Engineering Seminar Write-up

This web page summarizes the findings of a two-year UW Quantum System Engineering (QSE) Seminar, informally known as *Spinometers*.

Until recently this site was password-protected, and was open only to participants in the Spinometers Seminar. As of May 1, 2006 we are opening access to the general public.

Seminar review article (manuscript)

Here is the in-progress write-up of our Spinometers Seminar.

Version remarks

The most recent version is May 4, and the main upgrades are references to Serge Lang's *Algebra* textbook.

- [QSE is in NP.pdf](#): *Measurement operator methods for quantum model order reduction in large-scale spin simulations*
- [QSE is in NP.tar.gz](#): tarball of the LaTeX for the above

4 May 2006

Measurement operator methods for quantum model order reduction in large-scale spin simulations

J. A. Sidles¹†, D. D. Caplow², J. L. Garbini³, L. E. Harrell⁴, A. O. Hero⁵, J. P. Jacky¹, C. K. Kikuchi⁶, J. R. Malcomb³, A. G. Norman⁷

¹ Department of Orthopaedics and Sports Medicine, Box 356500, School of Medicine, University of Washington, Seattle, Washington, USA

² Department of Chemistry, College of Engineering, University of Washington, Seattle, Washington, USA

³ Department of Mechanical Engineering, College of Engineering, University of Washington, Seattle, Washington, USA

⁴ Department of Physics, U. S. Military Academy, West Point, New York, USA

⁵ Department of Electrical Engineering, College of Engineering, University of Michigan, Ann Arbor, Michigan, USA

⁶ Premedical Undergraduate Program, College of Arts and Sciences, University of Washington, Seattle, Washington, USA

⁷ Department of Bioengineering, College of Engineering, University of Washington, Seattle, Washington, USA

E-mail: sidles@u.washington.edu (John A. Sidles)

Abstract. This article introduces a new technique for model order reduction in large-scale quantum spin simulations. The technique has three stages: first the deliberate introduction of noise into the simulation; then the conversion of that noise into an equivalent continuous measurement process; and finally the projection of the resulting quantum trajectory onto a product-sum manifold of Beylkin-Mohlenkamp type. The application of this technique is illustrated by numerical quantum trajectory simulations of single-spin detection by magnetic resonance force microscopy (MRFM). Excellent agreement with experimental results is obtained. It is shown that in the Markovian noise limit single-spin data statistics can be predicted in closed form to all orders, despite the nonlinearity of the underlying quantum model. A new closed-form positive P -representation of the thermal density matrix is derived as an auxiliary result. Large-scale quantum simulations of up to eighteen interacting spins are then presented, in the context of a spin-dust model that has no symmetries, no spatial ordering, and high temperature; the technique is found to work well even within this deliberately challenging model. Classical and quantum model order reduction (MOR) techniques are compared, and the invariance associated with Choi's Theorem is shown to be essential to quantum model order reduction. A surprising equivalence between compressed wave functions and cryptographic keys is noted, and as a proof-of-principle, the numerical spin-dust trajectories are used to implement a public key exchange protocol. A large class of quantum systems can be simulated in polynomial space and time by this method, with the main restriction that the noise level be large enough that the system being simulated is not running a quantum computation.

PACS numbers: 82.20.Wt, 75.40.Gb, 03.65.Ud, 76.20.+q