

Quinone Induced Activation of Keap1/Nrf2 Signaling by Aspirin Prodrugs Masquerades as Nitric Oxide

TAREISHA DUNLAP, SUJEEWA C. PIYANKARAGE, GIHANI T. WIJEWICKRAMA, SAMER ABDUL-HAY,
MICHAEL VANNI, VLADISLAV LITOSH, JIA LUO, AND GREGORY R. J. THATCHER*

Department of Medicinal Chemistry & Pharmacognosy, University of Illinois College of Pharmacy,
University of Illinois at Chicago, 833 S Wood St, Chicago, IL 60612

*Correspondence: Gregory Thatcher, Department of Medicinal Chemistry and Pharmacognosy, University of
Illinois at Chicago, 833 S. Wood Street, Chicago, IL, 60612; E-mail: thatcher@uic.edu

Supporting Information

Figure S1: Effect of *p*BrASA on intracellular NO concentrations. DAF-DA fluorescence in HepG2 cells after 4 h treatment with DMSO, L-NAME (5 μ M), *p*Br-ASA (25 μ M) and L-NAME (5 μ M) after 30 min with *p*Br-ASA (25 μ M) after 4 h.




Figure S1: Effect of *p*BrASA on intracellular NO concentrations. DAF-DA fluorescence in HepG2 cells after 4 h treatment with DMSO, L-NAME (5 μ M), *p*Br-ASA (25 μ M) and L-NAME (5 μ M) after 30 min with *p*Br-ASA (25 μ M) after 4 h.