


1. Consider a sphere, of radius R and uniform charge density such that the total charge is Q .
 (a) (5) There is a second Gaussian sphere of radius $r > R$ with a center which is a distance $2r$ away from the center of the charge carrying sphere. Determine the electric flux through the Gaussian sphere.



- (b) (5) Now suppose the Gaussian sphere of radius $r > R$ has the same center as the original sphere. Determine the electric flux through the second sphere for this new situation.



- (c) (10) Consider again the Gaussian sphere of part (b). Show that the surface integral over the Gaussian sphere is given by $\oint \vec{E}(\vec{r}) \cdot \hat{n} dA = 4\pi r^2 E(r)$.

Spherically symmetric charge distribution

$$\vec{E}(\vec{r}) = E(r) \hat{r} \quad \text{Sphere } \hat{n} = \hat{r}$$

$$\vec{E} \cdot \hat{n} = E(r)$$

$$\oint_{\text{Sphere}} \vec{E} \cdot \hat{n} dA = \int_{\text{Sphere}} E(r) dA = E(r) \int dA = E(r) 4\pi r^2$$

- (d) (10) Consider $r < R$, show that $E(r) = \frac{1}{\epsilon_0} \frac{\rho Q}{4\pi R^3} r$.

$$E 4\pi r^2 = \text{charge enclosed} / \epsilon_0$$

$$= \frac{1}{\epsilon_0} \int_0^r 4\pi r'^2 dr' = \frac{Q}{4\pi R^3} \frac{4\pi r^3}{3}$$

$$\rho = \frac{Q}{4\pi R^3}$$

$$E 4\pi r^2 = \frac{1}{\epsilon_0} 4\pi \frac{r^3}{3} \frac{Q}{4\pi R^3}$$

$$\therefore E = \frac{1}{\epsilon_0} \frac{r^2 Q}{4\pi R^3}$$

(e) (15) Determine the electrostatic energy stored in the sphere of radius R and charge Q .

$$\begin{aligned} W &= \frac{1}{2} \epsilon_0 \int d^3r E^2 \\ &= \frac{\epsilon_0}{2} 4\pi \left\{ \int_0^R r^2 dr E^2 + \int_R^\infty r^2 dr E^2 \right\} \\ &= \epsilon_0 2\pi \left[\int_0^R r^2 dr \left(\frac{r-Q}{4\pi R^2 \epsilon_0} \right)^2 + \int_R^\infty r^2 dr \left(\frac{Q}{4\pi \epsilon_0 r^2} \right)^2 \right] \\ &= \frac{\epsilon_0 2\pi Q^2}{16\pi^2 \epsilon_0} \left[\int_0^R \frac{r^2 dr r^2}{R^6} + \int_R^\infty \frac{r^2 dr}{r^4} \right] \\ &= \frac{Q^2}{8\pi \epsilon_0} \left[\frac{R^5}{5R^6} + \frac{1}{R} \right] = \frac{Q^2}{4\pi \epsilon_0 R} \frac{3}{5} \end{aligned}$$

2. This problem is concerned with a thin disk of radius R with a charge per unit area of σ .

(a) (10) Compute the electric potential $V(z)$ at a distance z on the axis of the disk (directly above the center of the disk).

$V(P) = \frac{1}{4\pi\epsilon_0} \int \frac{da' \sigma}{|\vec{r} - \vec{r}'|} = \frac{1}{4\pi\epsilon_0} \sigma \int_0^R s ds \int_0^{2\pi} \frac{d\phi}{\sqrt{z^2 + s^2}}$
 $dq = \sigma da$
 $\int da' \sigma = \int_0^R \int_0^{2\pi} \sigma s ds d\phi = \frac{1}{4\pi\epsilon_0} \sigma 2\pi \int_0^R \frac{s ds}{\sqrt{s^2 + z^2}} = \frac{1}{4\pi\epsilon_0} 2\pi \sigma (\sqrt{R^2 + z^2} - z)$
 $da = s ds d\phi$


(b) (10) Determine the electric field $\vec{E}(z)$ at that same point.

$\vec{E} = E(z) \hat{z}$ (from cylindrical symmetry)
 $E(z) = -\frac{\partial V}{\partial z} = \frac{1}{2\epsilon_0} \left[\frac{z}{\sqrt{R^2 + z^2}} - 1 \right]$

(c) (10) Suppose $R \gg z$, what should the electric field $\vec{E}(z)$ be?

$R \gg z$ $E = \frac{\sigma}{2\epsilon_0}$ from above, or by application of Gauss Law

(d) (15) Now consider positions close to the axis, but a distance $s \ll R_3^c$ away from the axis. Obtain the potential $V(s, d)$. Your answer should be expressed in terms of a well-defined double integral. Do not carry out the integral.



$$V(s, d) = \frac{1}{4\pi\epsilon_0} \sigma \int_0^R s' ds' \int_0^{2\pi} d\phi \frac{1}{|\vec{r} - \vec{r}'|}$$

$$|\vec{r} - \vec{r}'| = \sqrt{z^2 + (x-x')^2 + (y-y')^2} = \sqrt{z^2 + s^2 + s'^2 - 2ss'\cos\phi}$$

$$V(s, d) = \frac{1}{4\pi\epsilon_0} \sigma \int_0^R s' ds' \int_0^{2\pi} \frac{d\phi}{\sqrt{z^2 + s^2 + s'^2 - 2ss'\cos\phi}}$$

(for $s \ll R, z$)

(e) (10) Explain why $V(s, z)$ is well approximated by the expression of the form $V(s, z) = V(0, z) + s^2 f(z)$.

$E_x = E_y = 0$ for $x = y = 0$, Cylindrical symmetry means $V(s, z)$ depends on s^2 .
Thus any dependence on s^2 must lead to vanishing E_x, E_y .

$$s^2 = x^2 + y^2$$

$$E_x \propto \frac{\partial V}{\partial x} = 0 \text{ for } x=0$$

$$E_y \propto \frac{\partial V}{\partial y} = 0 \text{ for } y=0$$

s^4 and higher powers are ignored because s^2 is small.