

Electrodynamics, Physics 323
Spring 2004

First midterm
 Instructor: David Cobden

8.20 am, April 23, 2004

You have 60 minutes. End on the buzzer at 9.20. Answer all 11 questions.

Write your name on every page and your ID on the first page.

Write all your working on these question sheets. Use this cover page for extra working (you might get credit for it.)


It is important to show your calculation or derivation. You won't get full marks just for stating the correct answer if you don't show how you get it.

Watch the blackboard for corrections or clarifications during the exam.

This is a **closed book** exam. **No notes allowed. No calculators.**

Do not turn this page until I say 'go'.

In a dilute plasma or in a waveguide, $\omega^2 = \omega_p^2 + c^2 k^2$



I. A plane electromagnetic wave, of electric field amplitude E_{0I} , is normally incident on a dielectric boundary across which the refractive index changes from n_1 to n_2 , where $n_1 > n_2$.

1. [5] What is the magnitude of the magnetic field in the incident wave, B_{0I} ?

2. [5] State the relationship between the incident, reflected and transmitted amplitudes, E_{0I} , E_{0R} and E_{0T} , resulting from the boundary condition on electric field.

3. [15] Assuming $\mu_1 = \mu_2$, the amplitude of the reflected wave is $E_{0R} = \frac{(n_1 - n_2)}{(n_1 + n_2)} E_{0I}$.

Using conservation of energy, or otherwise, derive the amplitude of the transmitted wave, E_{0T} .

II. Dilute plasma.

4. [10] Starting from the appropriate Maxwell equation(s) in a linear conducting medium, derive the dispersion relation: $k^2 = \mu\epsilon\omega^2 + i\sigma\mu\omega$.

5. [15] Show that electromagnetic waves cannot propagate in a dilute plasma for $\omega < \omega_p$, where the plasma frequency is $\omega_p = (Nq^2/m\epsilon_0)^{1/2}$. You should derive this result by considering a free charged particle oscillating in response to the electric field. Define all the terms in the equation for ω_p .

6. [10] At what speed, in terms of c , does a radio pulse of central frequency $\omega = 2\omega_p$ propagate through the plasma?

III. The inside of a (square) metal waveguide is defined by the region $0 < x < a$, $0 < y < a$. Consider only its *TM* modes.

7. [5] State B_z and show that it matches the boundary conditions on \mathbf{B} .

8. [10] We search for waveforms propagating in the positive z direction, of the form $\mathbf{E}(x, y)e^{i(kz - \omega t)}$. From Maxwell's wave equation for \mathbf{E} , derive a differential equation for $E_z(x, y)$ which involves ω but not time.

9. [5] State the boundary conditions on $E_z(x,y)$.

10. [15] Determine the set of functions $E_z(x,y)$, (corresponding to the TM_{mn} modes), that satisfy the equation from Q.8 and the boundary conditions from Q.9.

11. [5] Find the cutoff frequency for the lowest TM mode.