

# The time course and specificity of perceptual deterioration

Sara C. Mednick\*, A. Cyrus Arman, and Geoffrey M. Boynton

The Salk Institute for Biological Studies, SNL-B, 10010 North Torrey Pines Road, La Jolla, CA 92037-1099

Edited by Larry R. Squire, University of California, San Diego, CA, and approved January 24, 2005 (received for review October 28, 2004)

**Repeated within-day testing on a texture discrimination task leads to retinotopically specific decreases in performance. Although perceptual learning has been shown to be highly specific to the retinotopic location and characteristics of the trained stimulus, the specificity of perceptual deterioration has not been studied. We investigated the similarities between learning and deterioration by examining whether deterioration transfers to new distractor or target orientations or to the untrained eye. Participants performed a texture discrimination task in three one-hour sessions. We tested the specificity of deterioration in the final session by switching either the orientation of the background or the target elements by 90°. We found that performance deteriorated steadily both within and across the first two sessions and was specific to the target but not the distractor orientation. In a separate experiment, we found that deterioration transferred to the untrained eye. Changes in performance were independent of reported sleepiness and awareness of stimulus changes, arguing against the possibility that perceptual deterioration is due to general fatigue. Rather, we hypothesize that perceptual deterioration may be caused by changes in the ability for attention to selectively enhance the responses of relatively low-level orientation-selective sensory neurons, possibly within the primary visual cortex. Further, the differences in specificity profiles between learning and deterioration suggest separate underlying mechanisms that occur within the same cortical area.**

primary visual cortex | learning | vision | psychophysics | attention

It is said that practice makes perfect, but what happens with too much practice? Everyday experience suggests that intense training in a short period leads to a deterioration in performance, much like how muscles fatigue from lifting weights. In typical perception research, such overtraining is usually avoided rather than studied. This study, however, deliberately investigated how performance deteriorates throughout the day with repeated testing on a perceptual task.

Deterioration of performance with practice has been noted in prior studies. In the 1940s, Hull (1) reported increased reaction time with prolonged testing on simple repetitive tasks, such as letter cancellation, detecting differences in simple shapes, or adding three digits. This so-called “reactive inhibition” was defined as a tendency toward increased response times as a function of the number of repetitions and the intertrial interval. Hull proposed a multimodal fatigue factor that was driven by a natural and constantly increasing inclination to switch from the present task to another. Where reactive inhibition was a general principle, the present study examines deterioration as a specific mechanism of the visual system.

Although perceptual deterioration has received little scientific attention, there is a growing body of research on a related dynamical property of the visual system: perceptual learning. The perceptual learning literature provides a natural framework for studying perceptual deterioration because both learning and deterioration involve the effects of repeated testing on a task across sessions. Studies of perceptual learning show that for most tasks, learning does not transfer across stimulus properties such as retinotopic position (2), orientation (2, 3), spatial frequency

(4), motion direction and speed (5, 6) and, in some cases, even eye of origin (7) (for an excellent review, see ref. 8). This stimulus specificity suggests that perceptual learning may be mediated by neuronal plasticity in early, retinotopically organized visual areas. Stimulus specificities in perceptual learning, however, might also be influenced by neurons at higher stages of visual processing that become selective for properties such as orientation and spatial position as a consequence of training (9).

Mednick and colleagues (10, 11) found perceptual deterioration by using the same texture discrimination task developed to study learning (7). They reported that performance deteriorated significantly with each hour-long training session throughout the day. Importantly, this deterioration effect was retinotopically specific; when the target stimulus was shifted to the opposite visual hemifield for the final test session, performance returned to that of the first session. Such spatial specificity shows that perceptual deterioration is not simply due to general fatigue or boredom and further suggests that the effect can be attributed to plasticity of neurons in early visual cortex.

Given the initial similarities between learning and deterioration, we sought to determine whether, like learning, deterioration shows further specificity to the properties of the trained stimulus. In this study, we report that perceptual deterioration is binocular and specific to the target orientation, but not the background orientation. This pattern of effects is different from that of learning and suggests that deterioration does not necessarily share common neural mechanisms with learning.

Furthermore, prior studies using this task have only presented changes in thresholds across hour-long sessions rather than examining the buildup of deterioration within a session. This study investigates the development of deterioration both within and across sessions. We found that after a brief learning period within the first session, performance steadily decreases with continued exposure. This result shows that deterioration builds with the amount of exposure to the task across the day and not with the amount of time between training sessions (10).

## Methods

All experiments used the general procedure outlined in this section. Deviations from this procedure are described in each section.

**Participants and Procedures.** A total of 30 participants gave informed consent to participate in the study, which was approved by the internal review boards of both the University of California at San Diego and The Salk Institute for Biological Studies. All participants, ages 18–30 years, had normal or corrected-to-normal vision and no history of neurological, mental, or physical illness. Participants were restricted from caffeine the day of the study, restricted from alcohol the evening before test day, and were asked to get at least seven hours of sleep the night before

This paper was submitted directly (Track II) to the PNAS office.

Abbreviation: ISI, interstimulus interval.

\*To whom correspondence should be addressed. E-mail: smednick@salk.edu.

© 2005 by The National Academy of Sciences of the USA








Fig. 5. Transfer of deterioration in trained eye ( $T2 - T1$ ) and the untrained eye ( $T3 - T1$ ).

eye (236 vs. 319 ms, session one and two, respectively; paired  $t$  test,  $P = 0.009$ ). Difference scores were calculated to measure threshold ( $T$ ) changes in the same eye across sessions (Session 2( $T2$ ) – Session 1( $T1$ )) and in different eyes across sessions (Session 3( $T3$ ) – Session 1( $T1$ )). If deterioration were binocular, performance decreases in the trained eye would transfer to the untrained eye. No significant differences were found between thresholds obtained with the trained versus the untrained eye (paired  $t$  test,  $P = 0.39$ ) (mean and standard error shown in Fig. 5). Thus, deterioration appears to occur at least at the level of binocular neurons of early visual cortex.

### Conclusions

Herein, we have described the time course and specificity of perceptual deterioration that develops with repeated, within-day testing on a texture discrimination task. Deterioration occurred within the first testing session and continued to increase in a linear fashion with continued training. Thresholds remained dynamic throughout the entire testing period. An initial improvement, perhaps due to procedural (13) or perceptual (14) learning, was followed by a steady increase in subsequent thresholds. At no time during testing did performance remain stable. Thus, the concept of an average threshold becomes difficult to define for these data.

Previous studies using the texture discrimination task have shown performance improvement when participants were allowed a period of sleep between training and retesting (7, 11, 15). This sleep-dependent learning was specific to the eye of origin (7) (not replicated in ref. 16), the orientation of the background (but not the target) elements (7), and the retinotopic location of the target (7, 10), indicating slow, sleep-dependent plasticity of early visual cortex.

This study uses the same task as above but shows very different results. We report that unlike learning, deterioration is not specific to the eye of origin or the orientation of the background elements, but it is specific to the orientation of the target elements. These results indicate that deterioration is likely driven by rapid plasticity of early visual cortical areas. Thus, the processes pertaining to learning and deterioration appear to share the same brain areas but are mediated by different types of neuronal plasticity. This temporal difference in development (rapid for deterioration versus slow for learning), along with the difference in specificity (target versus background, binocular versus monocular), suggests two separate mechanisms underlying perceptual learning and deterioration, each sensitive to different visual information.

It has been suggested that the dependence of learning on the background orientation involves plasticity in contextual interactions between background and target (refs. 7 and 17; for a contrary view, see ref. 18). In contrast, deterioration appears

to develop solely for the attended target. Deterioration recovers when the target is moved to a new spatial location, even though oriented elements had been placed at this ignored location throughout previous training sessions (10). Thus, deterioration only seems to occur in mechanisms representing attended stimuli.

We noted a dissociation between subjective report and perceptual performance in the test of specificity. In agreement with behavioral data, most participants experienced the second session as harder than the first. In the third session, however, all participants were aware of the change in orientation of the background, but performance in this condition was no different from the control condition. On the other hand, not a single participant accurately noted the change in the target, even though the target change led to a complete restoration of performance to baseline levels. Three aspects of these data underscore the perceptual nature of these deterioration results and argue against a generalized fatigue principle proposed by Hull (1). First, an undetected change of the target components restored performance to baseline levels. Second, a detected, full visual field change of the background did not affect performance. Last, subjective sleepiness ratings did not increase significantly across the day.

The extent to which these unstable thresholds may have depended on the participant pool (mostly comprised of university undergraduates inexperienced with psychophysical testing and this particular task) is unknown. Studies of the dynamics of skill acquisition have reported differences on a variety of performance measures between naïve and expert subjects (19, 20). Future studies comparing experienced and naïve subjects on a similar deterioration-producing task can test the hypothesis that expertise may serve as a protection against rapid perceptual deterioration.

These results have implications for interpreting performance decreases reported by more applied areas of study. Investigations of the “useful visual field” (21), or the region of the visual field from that we can extract information during eye fixation, report degradation (i.e., driving simulator task shows response time slowing and decreased accuracy) under a variety of circumstances and individual characteristics (22, 23). Compared with the complex behavioral patterns required for driving, this study of performance deterioration at the perceptual level can be useful in describing what specific components of the driving task may be susceptible to deterioration. In this light, perceptual deterioration may have applications beyond being just a nuisance for psychophysicists planning an experimental design.

An analogous body of research to perceptual deterioration of the visual system can be found in sensory motor research showing practice-dependent performance decreases that are linked with cortical plasticity. It has been shown that even a small amount of tactile finger stimulation can produce rapid alteration of the orderly topographic digit representation of primary somatosensory cortex (24). Moreover, repetitive stereotyped hand movements have been shown to produce mild to severe motor deficits leading to acquired focal hand dystonia, associated with writer’s cramp, typist’s cramp, and musician’s cramp (25). fMRI studies have shown that focal hand dystonia has been correlated with altered somatotopic finger representations in humans (26). Further, physiological studies in monkeys have shown massive receptive field expansion, spreading of receptive fields from single digits to multiple digits, and dedifferentiation of the normally sharply segregated areas of the hand representation in area 3b (27, 28). It has also been noted that people with focal hand dystonia show impaired spatial and temporal tactile discrimination (29, 30). Thus, a repeated motor task can lead to performance decreases matched with distortions of neural representations.

Interestingly, intensive physical therapy of the affected hand can cause reorganization of somatotopic finger representations to normal borders and improve functioning of the affected hand (31), indicating a wide range of experience-dependent plasticity in the adult human cortex from learning to deterioration. These studies show that the experience-dependent neural plasticity underlying both learning and deterioration may be adaptive in terms of learning and maladaptive in terms of deterioration. An interesting speculation is that such neural plasticity may be occurring in visual cortex throughout the day.

Attention could lead to enhanced deterioration simply because attention enhances the response to the stimulus and, like adaptation, greater responses may cause more deterioration. Neuroimaging and physiological studies of attention both show that attention affects the magnitude of response as early as primary visual cortex (V1) (32–34). These attentional effects, however, are relatively small. Early visual areas still

respond well to unattended (35), and even unseen,<sup>†</sup> stimuli, whereas the unattended stimuli (background elements) in the present study had no effect on perceptual deterioration. Thus, it appears that attentional gain alone cannot explain the relationship between attention and deterioration.

Prolonged performance on a task may lead to deterioration in the ability to allocate attention to a stimulus attribute. Such a mechanism has been proposed for perceptual learning, wherein observers “learn to pay attention to the set of neurons that is most sensitive to the task” (36). Further, a recent neurophysiological study in the macaque shows that attention to a specific aspect of a stimulus selectively enhances the responses of V1 neurons in a very task-specific manner (37). Perhaps it is this task-specific mechanism of attention that is deteriorating over time.

<sup>†</sup>Whitney, D. G., Goltz, H. C. & Goodale, M.A. (2004) *J. Vis.* **4**, 44a (abstr.).

1. Hull, C. L. (1951) *Essentials of Behavior* (Greenwood, Westport, CT).
2. Schoups, A., Vogels, R. & Orban, G. (1995) *J. Physiol. (London)* **483**, 797–810.
3. Beard, B. L., Levi, D. M. & Reich, L. N. (1995) *Vision Res.* **35**, 1679–1690.
4. Fiorentini, A. & Berardi, N. (1980) *Nature* **287**, 43–44.
5. Liu, Z. & Vaina, L. M. (1998) *Brain Res. Cogn. Brain Res.* **6**, 347–349.
6. Matthews, N. & Welch, L. (1997) *Percept. Psychophys.* **59**, 60–72.
7. Karni, A. & Sagi, D. (1991) *Proc. Natl. Acad. Sci. USA* **88**, 4966–4970.
8. Fine, I. & Jacobs, R. (2002) *J. Vis.* **2**, 190–203.
9. Mollon, J. D. & Danilova, M. V. (1996) *Spat. Vis.* **10**, 51–58.
10. Mednick, S. C., Nakayama, K., Cantero, J. L., Atienza, M., Levin, A. A., Pathak, N. & Stickgold, R. (2002) *Nat. Neurosci.* **5**, 677–681.
11. Mednick, S., Nakayama, K. & Stickgold, R. (2003) *Nat. Neurosci.* **6**, 697–698.
12. Pelli, D. G. (1997) *Spat. Vis.* **10**, 437–442.
13. Karni, A. & Sagi, D. (1993) *Nature* **365**, 250–252.
14. Hawkey, D. J., Amitay, S. & Moore, D. R. (2004) *Nat. Neurosci.* **7**, 1055–1056.
15. Stickgold, R., James, L. & Hobson, J. A. (2000) *Nat. Neurosci.* **3**, 1237–1238.
16. Schoups, A. A. & Orban, G. A. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 7358–7362.
17. Adini, Y., Sagi, D. & Tsodyks, M. (2002) *Nature* **415**, 790–793.
18. Yu, C., Klein, S. A. & Levi, D. M. (2004) *J. Vis.* **4**, 169–182.
19. Milton, J. G., Small, S. S. & Solodkin, A. (2004) *J. Clin. Neurophysiol.* **21**, 134–143.
20. Gray, R. (2004) *J. Exp. Psychol. Appl.* **10**, 42–54.
21. Mackworth, J. F. (1970) *Vigilance and Attention: A Signal Detection Approach* (Penguin Books, Baltimore).
22. Roge, J., Pebayle, T., Lambilliotte, E., Spitzstetter, F., Giselbrecht, D. & Muzet, A. (2004) *Vision Res.* **44**, 2737–2744.
23. Roge, J., Pebayle, T., El Hannachi, S. & Muzet, A. (2003) *Vision Res.* **43**, 1465–1472.
24. Ziemus, B., Huonker, R., Haueisen, J., Liepert, J., Spengler, F. & Weiller, C. (2000) *NeuroReport* **11**, 1285–1288.
25. Nudo, R. J. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 7425–7427.
26. Elbert, T., Candia, V., Altenmuller, E., Rau, H., Sterr, A., Rockstroh, B., Pantev, C. & Taub, E. (1998) *NeuroReport* **9**, 3571–3575.
27. Jenkins, W. M., Merzenich, M. M. & Recanzone, G. (1990) *Neuropsychologia* **28**, 573–584.
28. Byl, N. N., Merzenich, M. M. & Jenkins, W. M. (1996) *Neurology* **47**, 508–520.
29. Molloy, F. M., Carr, T. D., Zeuner, K. E., Dambrosia, J. M. & Hallett, M. (2003) *Brain* **126**, 2175–2182.
30. Sanger, T. D., Tarsy, D. & Pascual-Leone, A. (2001) *Movement Disorders* **16**, 94–99.
31. Candia, V., Wienbruch, C., Elbert, T., Rockstroh, B. & Ray, W. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 7942–7946.
32. Gandhi, S. P., Heeger, D. J. & Boynton, G. M. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 3314–3319.
33. Martinez, A., Anillo-Vento, L., Sereno, M. I., Frank, L. R., Buxton, R. B., Dubowitz, D. J., Wong, E. C., Hinrichs, H., Heinze, H. J. & Hillyard, S. A. (1999) *Nat. Neurosci.* **2**, 364–369.
34. Somers, D. C., Dale, A. M., Seiffert, A. E. & Tootell, R. B. H. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 1663–1668.
35. Rees, G., Russell, C., Frith, C. D. & Driver, J. (1999) *Science* **286**, 2504–2507.
36. Saarinen, J. & Levi, D. M. (1995) *Vision Res.* **35**, 519–527.
37. Li, W., Piech, V. & Gilbert, C. D. (2004) *Nat. Neurosci.* **7**, 651–657.