Plant Propagation Protocol for *Grindelia stricta* ESRM 412 – Native Plant Production

	TAXONOMY
Plant Family	
Scientific Name	Asteraceae
Common Name	Daisy
Species Scientific Name	
Scientific Name	
Genus:	Grindelia
Species:	stricta
Species Authority:	DC.
Varieties	
Sub-species	
Cultivar	
Common Synonym(s)	Grindelia integrifolia DC.
January Car	Grindelia integrifolia var. macrophylla
	Grindelia macrophylla
Common Name(s)	Oregon gumweed,
Species Code (as per USDA Plants	GRST3
database)	
GENERAL INFORMATION	
Geographical range	Oregon gumweed is perennial herbaceous species found in North America from southern BC to northern California where it is found in beaches, rocky bluffs, and disturbed sites. Flowers are yellow, daisy-like; it is summer flowering. Appearance varies with the environment
Ecological distribution	Mounding, spreading groundcover especially suited for coastal sites, yellow, daisy-like flowers in summer contrast with ample green foliage. They need dry soil and full sun, and are raised from seed. They are not

	adapted to regions of cold, wet winters.	
Climate and elevation range	0 - 1640 feet	
Local habitat and abundance	0 - 1040 1001	
Local habitat and abundance		
Plant strategy type / successional	Strong stress-toleration. Sun, drought tolerant or	
stage	occasional summer water, adaptable to many kinds of	
	soil. This is a highly variable species which has	
	received various treatments. Douglas (1995) concluded	
	that it was best to recognize only one environmentally	
Di di di di di	variable species.	
Plant characteristics	Herb to 90 cm; stems simple or branched, decumbent	
	to ascending, glabrous to villous, occasionally	
	glandular. Lvs to 25*4cm, oblong to oblanceolate, acute to obtuse, entire to minutely serrate near apex,	
	base attenuate, petiolate; stem lvs amplexicaul.	
	base attenuate, petiorate, stem ivs ampiexicaur.	
PROPAGATION DETAILS		
Ecotype	Colorado	
Propagation Goal	Seed.	
Propagation Method	Seed.	
Product Type	Propagules (seeds, cuttings, poles, etc.)	
Stock Type		
Time to Grow	1 week.	
Target Specifications	80 cm	
Propagule Collection Instructions	Collect seeds in the fall from flowering heads	
Propagule Processing/Propagule	140 per gram.	
Characteristics	Sand commington in 2.2 weeks at 50.60 % Saw in shown	
Pre-Planting Propagule Treatments	Seed germinates in 2-3 weeks at 50-60 °F. Sow in sharp draining seed mix and cover to depth of seed	
Growing Area Preparation / Annual		
Practices for Perennial Crops		
Establishment Phase Details		
Length of Establishment Phase	Time to germination: 6 days.	
Active Growth Phase	2 month.	
Length of Active Growth Phase		
Hardening Phase		
Length of Hardening Phase		
Harvesting, Storage and Shipping	Seed storage condition: Seed stored in the greenhouse.	
Length of Storage		
Guidelines for Outplanting /		
Performance on Typical Sites		
Other Comments INFORMATION SOURCES		
INFORMATION SOURCES		

References	See below
Other Sources Consulted	
Protocol Author	Chao Yang
Date Protocol Created or Updated	05/19/2015
-	

Butler, Jennifer; Frieswyk, Christin. 2001. Propagation protocol for production of Grindelia seeds; USDI NPS - Rocky Mountain National Park, Estes Park, Colorado. In: Native Plant Network. URL: http://www.nativeplantnetwork.org (accessed 19 May 2015). Moscow (ID): University of Idaho, College of Natural Resources, Forest Research Nursery.

Bauer, Nancy. *The California Wildlife Habitat Garden: How to Attract Bees, Butterflies, Birds, and Other Animals*. Berkeley, CA: U of California, 2012. 198-199. *Google Book*. Web. 18 May 2015.

Everett, Thomas H. The New York Botanical Garden Illustrated Encyclopedia of Horticulture. P1557. New York: Garland Pub., 1980. Print.

Huxley, Anthony Julian, and Mark Griffiths. Dictionary of Gardening. London: Macmillan, 1992. Print.

McVicar, Jekka. Seeds: The Ultimate Guide to Growing Sucessfully from Seed. Conn.: Lyons, 2003. Print.

Ravetta, Damian A. "Download PDFs." Resin Production of Grindelia Accessions under Cultivation. N.p., n.d. Web. 20 May 2015.

http://www.sciencedirect.com/science/article/pii/0926669096894490.

Olivieri, Alejandro C. "Stereoselective Synthesis of the Novel Bisnorditerpene Grindelistrictic Acid, Isolated from Grindelia Stricta." - The Journal of Organic Chemistry (ACS Publications). N.p., July 1986. Web. 20 May 2015. http://pubs.acs.org/doi/abs/10.1021/jo00364a043>.

Goman, Michelle. "Statistical Analysis of Modern Seed Assemblages from the San Francisco." Bay: Applications for the Reconstruction of Paleo-salinity and Paleo-tidal Inundation. N.p., 01 Nov. 2001. Web. 20 May 2015. http://link.springer.com/article/10.1023/A:1012687113269>.

Dunford, Max P. "A Cytogenetic Analysis of Certain Polyploids in Grindelia (Compositae)." American Journal of Botany 51.1 (1964): 49-56. JSTOR. Web. 20 May 2015. http://www.jstor.org/stable/10.2307/2440063?ref=no-x-route:6aa20025f42b056198cef43efa4ff331.

Wayne, Lisa Beth. Recruitment Response to Salinity in Grindelia Stricta Var. Angustifolia: A Potential Indicator Species. N.p.: n.p., 1995. Print.

"Grindelia Stricta." Wikipedia. Wikimedia Foundation, n.d. Web. 20 May 2015. http://en.wikipedia.org/wiki/Grindelia_stricta. http://plants.usda.gov/core/profile?symbol=grst3