Roadmap

- Chromosome evolution:
 - Inversion
 - Transposition
 - Fission and fusion
- If today's lecture is perplexing, there is a useful film at http://www.youtube.com/watch?v=ZcnyMMHLkAw

One-minute responses

- Q: What is the difference between subfunctionalization and specialization?
 - Subfunctionalization: the gene has two functions, and the two new copies split them
 - Specialization: the gene has one function, but the two new copies adapt to do it in different tissues, times, or situations

One-minute responses

- Q: Are blue receptors and rhodopsins related to R/G and part of a larger gene family?
 - Yes. But those duplications are much older and not tandem.
- Q: How about olfactory receptors?
 - Yes, a HUGE superfamily with some human-specific branches (why?)





Figure 22. Phylogeny of vertebrate cone and rod opsins

An opsin mystery

- Dragonfish live in the deep sea, where most fishes see only blue light
- Dead specimens proved to have members of the red-receptor gene family—why?
- As soon as deep-sea explorers saw a live one, they understood....


Chromosomes

- Bacterial chromosomes are circular
- Most higher organisms have linear chromosomes with a centromere that attaches them to the spindle
- Telomeres at the ends protect the DNA from unravelling




An exception to the rule

- C. elegans, the lab nematode, has holocentric chromosomes
- The normal function of the centromere is diffused across the whole chromosome
- How did this evolve? Why did it evolve?



Chromosomes

 Through crossing-over chromosome strands can mix and recombine




 If gene order differs on the homologs this process gets into trouble

Chromosome rearrangements

- Inversion (changing the direction of part of the chromosome)
- Translocation (exchanging material between different chromosomes)
- Transposition (moving material to another location on the same chromosome)

Polytene chromosomes




- In flies and mosquitoes, "giant chromosomes" in salivary glands
- Chromosomes pair and then reduplicate themselves up to 1024 copies

Inversions

Paracentric inversion (does not include centromere)




Pericentric inversion (includes centromere)




I prefer not to use these words as they sound too similar to me!


Paracentric inversion heterozygote




Paracentric inversion heterozygote




Pairing in a paracentric inversion heterozygote






Crossing-over in a paracentric inversion heterozygote



Crossing-over in a paracentric inversion heterozygote



Paracentric inversion heterozygote - Outcome

We produce one of each:

Normal chromosome


Inverted chromosome

No centromere


$$A-B-C-D-E-F-G-B-A$$

Two centromeres (breaks randomly)


Pairing in a pericentric inversion heterozygote



Crossing-over in a pericentric inversion heterozygote



Crossing-over in a pericentric inversion heterozygote



Pericentric inversion heterozygote - Outcome

We produce one of each:

Normal chromosome

Inverted chromosome

Duplicate ABC, delete IJ


Duplicate IJ, delete ABC

Inversion summary

- Paracentric (doesn't include centromere)
 - Recombination produces two-centromere and no-centromere chromosomes
 - Recombinant gametes almost always die
- Pericentric (includes centromere)
 - Recombination produces duplications and deletions
 - Recombinant gametes might be viable if affected area is small
- If there is only one recombination, two of the four chromatids will be okay (but non-recombinant)

Evolutionary consequences of inversion

- Inversions "suppress recombination" really they kill recombinants
 - Reduced fertility bad
 - Preserve favorable groupings of alleles possibly good
- In most species, fixing an inversion requires drift or strong positive selection
- Some species have genetic systems more permissive of inversions




(c) Steven M. Carr

Multiple inversions have happened between different lineages of Drosophila

Drosophila inversion clines

- $Drosophila\ subobscura$ shows many inversions in the south relative to the north
- This cline was recreated in Chile and the West Coast of the USA after introduction of flies from Spain.



Drosophila inversion clines

- Inversions are unusually common in Drosophila why?
 - No recombination in males
 - Female oogenesis preferentially uses a cell with no broken chromosomes
 - Populations often established by a single female strong genetic drift

Practice problem

What, if any, problems arise for:

- A male Drosophila inversion heterozygote?
- A female Drosophila inversion heterozygote?
- A human inversion heterozogyote?
- A human inversion *homozogyote*?

Assume no genes were damaged by creation of the inversion

Practice problem

What, if any, problems arise for:

- A male Drosophila inversion heterozygote? Should be fine
- A female Drosophila inversion heterozygote? *If she has* multiple crossovers in the inversion, she may not make a viable egg, so she will have some fertility reduction
- A human inversion heterozogyote? Fertility reduction
- A human inversion homozogyote? Should be fine


Assume no genes were damaged by creation of the inversion

Translocation




A translocation

Before



A translocation heterozygote

at first division of meiosis metaphase




A pair of translocated chromosomes pairs with a pair of untranslocated chromosomes

Translocation

- Translocation arises from:
 - Breakage and repair
 - Illegitimate recombination between different chromosomes
- Translocation is usually reciprocal because every chromosome end must have a telomere
- Even without recombination, translocation heterozygotes mis-segregate their chromosomes in meiosis
- Translocations almost always underdominant
- Only drift or strong selection can allow one to spread

Why is translocation a problem?

- The strange pairing is fine in itself
- Recombination is fine
- Separation of the chromosomes in meiosis can have two outcomes:
 - Two translocated chromosomes go to one daughter and two normal to the other – fine
 - Each daughter gets one translocated and one not not good
 - This is a 50/50 chance



Practice problem

What, if any, problems arise for:

- A male Drosophila translocation heterozygote?
- A female Drosophila translocation heterozygote?
- A human translocation heterozogyote?
- A human translocation *homozogyote*?

Practice problem

What, if any, problems arise for:


- A male Drosophila translocation heterozygote? Fertility reduction due to mis-segregation
- A female Drosophila translocation heterozygote? Fertility reduction due to mis-segregation
- A human translocation heterozogyote? Fertility reduction due to mis-segregation; this is one cause of Down's Syndrome
- A human translocation homozogyote? Should be fine

Chromosome fission and fusion


Chromosome fusion probably explains why domestic horses have 64 chromosomes. . .



... and the closest living wild species and probable ancestor of domestic horses, <u>Przewalski's horse</u>, has 66 chromosomes.



Chinese muntjac deer Muntiacus reevesi



Indian muntjac deer Muntiacus muntjak Translocation, chromosome fusion, and/or fission explain why these two very similar species of hoofed mammal, the Chinese and Indian muntjac deer, have such different karyotypes.

Chromosome fission and fusion

- Fissions and fusions may begin as translocations where the reciprocal product is lost
- Generally underdominant
- This contributes to the sterility of mules (hybrids between two closely related species which differ in chromosome number)
- Closely related species often differ in chromosome number
- This may represent an early step in reproductive isolation

One-minute responses

- Tear off a half-sheet of paper
- Write one line about the lecture:
 - Was anything unclear?
 - Did anything work particularly well?
 - What could be better?
- Leave at the back on your way out