


Now that we have looked at the *physical* processes involved with the exchange of CO₂ between the atmosphere and the ocean let's turn to the *chemical* processes



IPCC 2007 Fig. 7.10

Chemical Processes Influencing Air-Sea Exchange of CO₂

1. Physical Processes

- Air-sea gas exchange = f (wind speed, bubble injection, surfactants)
- Ocean circulation


2. Chemical Processes

- CO₂ solubility = f (temperature, salinity) ["The Solubility Pump"]
- Carbonate chemical equilibrium

3. Biological Processes ["The Biological Pump"]

- Photosynthesis & respiration
- Calcium carbonate production

Solubility of Gases as a Function of Temperature




- Solubility of all gases *decreases with increasing T*
- Differences result from molecular interactions between gas & water

Sarmiento and Gruber (2006)

CO_2 Solubility is a Function of Temperature

- Demonstrations of the temperature dependence




- Solubility decreases as T increases


- $p\text{CO}_2$ increases as T increases (implying the liquid can hold less of the gas at higher T, resulting in higher gas pressure)

McGillis & Wanninkhof (2006) Aqueous CO_2 gradients for air-sea flux estimates, Mar. Chem. Vol. 98: 100-108

Gas Solubility Decreases as Salinity Increases



The Carbon "Solubility Pump"



- CO₂ is more soluble in cold waters than in warm waters (the **thermal pump**), & more soluble in fresher waters than saltier waters (the **salt pump**).
- If alkalinity (see following discussion) were uniform throughout the ocean & if both cold & warm surface waters equilibrated their pCO₂ with the atmosphere, then cold surface waters would have a higher dissolved CO₂ content than warm surface waters.
- As these cold surface waters circulate into the deep interior of the ocean, deep waters will have more CO₂ than warm surface waters.

Chemical Processes Influencing Air-Sea Exchange of CO₂

1. Physical Processes

- Air-sea gas exchange = f (wind speed, bubble injection, surfactants)
- Ocean circulation

2. Chemical Processes

- CO₂ solubility = f (temperature, salinity) ["The Solubility Pump"]
- Carbonate chemical equilibrium

3. Biological Processes ["The Biological Pump"]

- Photosynthesis & respiration
- Calcium carbonate production

CO₂ Uptake by the Ocean: the Carbonate System

- The ocean can take up CO₂ from the atmosphere in amounts that far exceed what would be expected based on solubility alone
- The extra absorbing power is caused by the **carbonate buffer system**
 - CO₂ gas dissolves in seawater to become aqueous CO₂

$$\text{CO}_2(\text{g}) \leftrightarrow \text{CO}_2(\text{aq})$$
 - *Henry's Law* describes the equilibrium between CO₂(g) & CO₂(aq)

$$K_H = [\text{CO}_2(\text{aq})]/p\text{CO}_2 \quad (\text{a function of T \& S})$$
 - CO₂(aq) combines with water to produce **carbonic acid**


$$\text{CO}_2(\text{aq}) + \text{H}_2\text{O} \leftrightarrow \text{H}_2\text{CO}_3$$
 - At the pH of surface seawater (~8.2), **carbonic acid** rapidly dissociates into a hydrogen ion and a **bicarbonate** ion

$$\text{H}_2\text{CO}_3 \leftrightarrow \text{H}^+ + \text{HCO}_3^-$$
 - The hydrogen ion then reacts with a **carbonate** ion to produce a second **bicarbonate** ion


$$\text{H}^+ + \text{CO}_3^{2-} \leftrightarrow \text{HCO}_3^-$$

CO₂ Uptake by the Ocean: DIC

- Since only ~1% of CO₂(aq) exists as H₂CO₃ it is usually left out of the sum of dissolved inorganic carbon (DIC) species




- The molar ratio of these three species in seawater is about 1 : 100 : 10



CO₂ Uptake by the Ocean: Conservative Quantities

- DIC is a **conservative** quantity in seawater, meaning
 - Its concentration can only be changed by mixing & advection
 - It can be mixed linearly
 - Non-conservative properties (e.g., O₂ & PO₄³⁻) are altered by biological & chemical processes

- An example would be mixing in an estuary, where the DIC concentration changes linearly from the low value in rivers to the high value in seawater:




*** Start Here Tues 2/10/09 ***

Clarifications

Eddies


- Small-scale ($\sim 10^0$ - 10^2 km), time-varying components of the circulation
- Spatio-temporal integrations produce the large-scale “steady” flows
- Contain 90-99% of the kinetic energy of the flow (Wunsch, 2004, 2007, in press)
- Ubiquity & importance recognized since '70s; widely observed since '90s



CO₂ Uptake by the Ocean: Alkalinity

- Another conservative quantity in seawater that is important for understanding the exchange of CO₂ between the atmosphere & the ocean is **Alkalinity**, the best definition of which I have ever read is:

The negative charge deficit in seawater that is compensated by ions which can exist in more than one charge state. (Broecker, 2005)



- Seawater must be **electrically neutral**
- Though comprised primarily of ions with a fixed electrical charge, such as the **cations** Na⁺, K⁺, Mg²⁺, Ca²⁺, and the **anions** Cl⁻ & SO₄²⁻, there is a *slight* deficit of negative charge, the **alkalinity**
- That deficit is made up by **protonating & deprotonating acids & bases** until charge balance is achieved

CO₂ Uptake by the Ocean: Alkalinity (cont'd.)

- pH is a 3rd important quantity in determining ocean uptake of CO₂

$$\text{pH} = -\log[\text{H}^+]$$

- The NCD is compensated by adding & removing protons (H⁺) to ions that can exist in more than one charge state
- Important for ocean CO₂ uptake are acids & bases that can exist in more than one charge state in the pH range of seawater, ~7-9.
- So a more precise definition of alkalinity would be:


"The negative charge deficit that is compensated by acids & bases that can exist in more than one charge state in the pH range of seawater."

- Because of their high concentration in seawater and their tendency to protonate / deprotonate at pH 7-9, HCO₃⁻, CO₃²⁻, B(OH)₃, & B(OH)₄⁻ are by far the most abundant such species

Adapted from Broecker (2005), Emerson & Hedges (2007) Chap. 4.

CO₂ Uptake by the Ocean: Alkalinity (cont'd)

- The high concentrations of carbonate & borate ions make HCO₃⁻, CO₃²⁻, B(OH)₃, & B(OH)₄⁻ the most important contributors to alkalinity in seawater since its pH is ~8.2



Adapted from Emerson & Hedges (2007) Fig. 4.2

CO₂ Uptake by the Ocean: Alkalinity (cont'd.)

PROPERTIES	TROPICAL SURFACE	DEEP PACIFIC	UNITS
WATER DEPTH	0	4000	meters
TEMPERATURE	25.0	1.5	°C
SALINITY	35.0	34.7	g/kg
P _{CO₂}	280	510*	10 ⁻⁶ atm
NCD	2216.0	2450.0	10 ⁻⁶ mol/kg
ΣCO ₂	1858.0	2340.0	10 ⁻⁶ mol/kg
(CO ₂) _{aq}	7.9	29.9	10 ⁻⁶ mol/kg
(HCO ₃ ⁻)	1601.5	2215.0	10 ⁻⁶ mol/kg
(CO ₃ ²⁻)	248.5	95.0	10 ⁻⁶ mol/kg
ΣB	410.6	407.1	10 ⁻⁶ mol/kg
B(OH) ₃ ⁰	302.0	362.6	10 ⁻⁶ mol/kg
B(OH) ₄ ⁻	108.6	44.5	10 ⁻⁶ mol/kg
(OH ⁻)	8.8	0.4	10 ⁻⁶ mol/kg
pH	8.15	7.77	

Borate contribution to Alkalinity:
 $108.6 / 2216 * 100 = 4.9\%$

Adapted from Broecker (2005) and Emerson & Hedges (2007) Chap. 4

- Thus in seawater:

$$\text{NCD} = \text{Alkalinity} = \text{HCO}_3^- + 2*\text{CO}_3^{2-} + \text{B(OH)}_4^-$$



- More precise definitions of **alkalinity** exist that include more species
- They include ions that contribute < 1% to alkalinity b/c they become protonated / deprotonated at:
 - seawater pH, but have low &/or variable *concentrations* (e.g., H_3SiO_4^- , H_2PO_4^- , HPO_4^{2-} , PO_4^{3-} , OH^-)
 - pH levels << 8 (e.g., Cl^- , SO_4^{2-} , F^-)
- In practice, even **borate** is left out since it comprises < 5% of alkalinity
- The term **Carbonate Alkalinity** is then used as a simplifying approximation:

$$\text{Carbonate Alkalinity} = \text{HCO}_3^- + 2*\text{CO}_3^{2-}$$

How is Alkalinity Determined?


- Alkalinity is measured by **titration**: strong acid is added to seawater until all proton acceptors have been protonated
- "The precise definition of alkalinity of seawater is based on the method by which it is determined and the species that exchange protons during the titration." (Emerson & Hedges (2007) p. 4.10)
- Operationally **carbonate alkalinity** is defined as the # of equivalents of acid required to bring a sample to the **CO₂ endpoint, or equivalence point** —i.e., when # moles acid added = moles HCO₃⁻ + 2*moles of CO₃²⁻

- Acid dropped from a burette into seawater until indicator changes color permanently--the endpoint.



- At this **equivalence point** all the CO₃²⁻ & HCO₃⁻ have been converted to CO₂
- The **carbonate alkalinity** is 3.8 meq/L


Alkalinity Titration of Seawater with Strong Acid



Zeebe & Wolf-Gladrow (2001) Fig. 1.2.8

Alkalinity in the Ocean

- Alkalinity is greater in the deep ocean than in the surface b/c Ca^{2+} ions are incorporated into CaCO_3 (removing alkalinity & NCD) in surface waters & released by dissolution of CaCO_3 in deep waters (adding alkalinity & NCD)



Alkalinity
(μMol or $\mu\text{eq/kg}$)

WOCE Pacific Ocean
Line 18 along 105°W

Adapted from Broecker (2005). WOCE Pacific Ocean Atlas.

*A biological process we will discuss shortly

CO₂ Uptake by the Ocean: the Revelle Factor





Figure adapted from Ed Boyle 12.842 Lecture Notes (2008)

- The amount of CO₂ the ocean can absorb from the atmosphere beyond the amount of CO₂ gas that can be dissolved in it is referred to as its **buffering capacity**
- The buffering capacity of the ocean is quantified by the **Revelle Factor**
- The **RF (aka buffer factor)** relates the fractional change in atmospheric pCO₂ to a fractional change in DIC (after re-equilibration)
- **RF** is directly proportional to the ratio of DIC : Alkalinity (see next page...)

Buffering Capacity of the Ocean


- Because **Carbonate Alkalinity** = HCO₃⁻ + 2*CO₃²⁻ it is clear that the greater the alkalinity of a solution the greater its potential for neutralizing acid (H⁺), such as CO₂ :



- The more CO₃²⁻, or **Alkalinity**, contained in the water, the more CO₂ (acid) it will be able to absorb
- But the total amount of DIC is also a factor, because with low concentrations of DIC there cannot be large amounts of CO₃²⁻ to neutralize CO₂ at any pH
- That is why the **Revelle factor** is proportional to (DIC / Alkalinity). Low values of **RF** of DIC/Alk imply greater capacity to buffer added CO₂. (Higher **Alk** = Lower **RF** = Higher buffering capacity)

Buffering Capacity of the Ocean: The Revelle Factor

1994 distribution of the **Revelle factor** averaged over upper 50m of water.




• Lower **RF** = Greater buffering capacity

- Low RFs occur in warm tropical-subtropical waters; high RFs in cold high latitude waters
- The capacity for seawater to take up CO_2 from the atmosphere is inversely proportional to the RF
- Hence, the lower the RF, the higher the oceanic equilibrium concentration of CO_2 for a given atmospheric CO_2 increase
- North Pacific surface waters have a higher RF at comparable latitudes & consequently lower anthropogenic CO_2 concentrations
- This difference results from North Pacific alkalinity values about 100 mol/kg lower than in the North Atlantic
- Current RFs are about one unit higher than in the preindustrial ocean.

Sabine et al. (2004) *Science* Vol. 305: 367-371

Equilibration Time for Atmospheric Gases in Ocean Mixed Layer


Simple Gas Exchange
(e.g. O_2 , Ar)



- For mixed layer depth of 100m & piston velocity of 2000 m/yr
- Equilibration time
 $= 100\text{m} / 2000\text{m yr}^{-1}$
 $= 0.05\text{yr} = 18\text{d}$

Equilibration Time for Atmospheric Gases in the Ocean Mixed Layer

pCO₂ equilibration




- Change in pCO₂ causes equal change in CO₂(aq)
- Revelle factor: fractional change in pCO₂ is ~10x greater than DIC change
- Since CO₂(aq)=0.5% of DIC it takes 10/0.5, or 20x longer to equilibrate DIC than to equilibrate CO₂(aq)
- Equilibration time

$$= (100\text{m} / 2000\text{m yr}^{-1}) * 20 = 1\text{yr}$$

Equilibration Time for Atmospheric Gases in the Ocean Mixed Layer

carbon isotope equilibration
(C13, C14)



$$2000 \text{ m yr}^{-1} * 10^{-5} \text{ moles kg}^{-1} * 1000 \text{ kg m}^{-3} = 20 \text{ moles m}^{-2} \text{ yr}^{-1}$$


Piston velocity	conc. of gaseous dissolved CO ₂	conversion Factor for H ₂ O	Exchange rate of CO ₂ across air/sea interface
--------------------	--	--	--

~[DIC] in seawater

- Total DIC must equilibrate with atm. CO₂
- 1m² of upper 100m of ocean contains: 10⁵kg water * (2000*10⁻⁶mol C/kg water) = 200 moles C
- CO₂ gas exchange rate = 20 mol/m²/yr (see prev. lec.)
- Equilibration time

$$= (200 \text{ mol}) / (20 \text{ mol/m}^2/\text{yr}) = 10\text{yr}$$

Now that we have looked at the *physical* & *chemical* processes involved with the exchange of CO_2 between the atmosphere & the ocean let's turn to the *biological* processes



IPCC 2007 Fig. 7.10

Biological Processes Influencing Air-Sea Exchange of CO_2

1. Physical Processes

- Air-sea gas exchange = f (wind speed, bubble injection, surfactants)
- Ocean circulation

2. Chemical Processes


- CO_2 solubility = f (temperature, salinity) ["The Solubility Pump"]
- Carbonate chemical equilibrium

3. Biological Processes ["The Biological Pump"]

- Photosynthesis & respiration
- Calcium carbonate production


The “Biological Carbon Pump”

- The biologically mediated transfer of CO_2 & nutrients from the surface to the deep ocean



* Redfield, Ketchum & Richards (1963) “The influence of organisms on the composition of sea-water.” In: M.N. Hill (Ed.), *The Sea* 2: 26-77, Interscience, NY.


A More Realistic View of Biological Processes in the Surface Ocean “Box”



POM = Particulate Organic Matter
DOM = Dissolved Organic Matter

Emerson & Hedges (2007) *Chem. Oceanogr.*, Fig. 6.5



Biological Pump Effect on Nutrient Distributions



- Nutrients such as N & P stripped out of surface water by **photosynthesis**
 - Consuming $\text{CO}_2(\text{aq})$
 - Producing O_2
- Transported to deep sea by **sinking fecal pellets**
- Released back into deep water during decomposition by bacteria (**rem mineralization**)
 - Consuming O_2

Shaffer (1996) *J. Geophys. Res.* Vol. 101(C2): 3723-3745.


Biological Pump Effect on Nutrient Distributions



- Note that O_2 has mirror-image profile of PO_4^{3-}


Shaffer (1996) *J. Geophys. Res.* Vol. 101(C2): 3723-3745. GEOSECS data.

Combined Effect of the Biological Pump & Ocean Circulation



PO4: Shaffer (1996) *J. Geophys. Res.* Vol. 101(C2): 3723-3745. MOC: John Marshall, MIT.


Biological Pump + Ocean Circulation Effect on Phosphate



- Oldest water in ocean accumulates most remineralized PO_4^{3-} , the ultimate source of which was photosynthesis in the global surface ocean.

WOCE (2007) Atlas Volume 2: Pacific Ocean


Biological Pump + Ocean Circulation Effect on DIC



- Oldest water in ocean accumulates most respired CO₂, the ultimate source of which was photosynthesis in the global surface ocean.

WOCE (2007) Atlas Volume 2: Pacific Ocean


Biological Pump + Ocean Circulation Effect on Oxygen



- Oldest water in ocean is most impoverished in O₂ because 138 moles of O₂ are consumed per mole of PO₄³⁻ liberated during organic matter decomposition
- From the Redfield ratio of C:N:P:O₂ of 106:16:1:138

WOCE (2007) Atlas Volume 2: Pacific Ocean

Global Nitrate & Water Age at 3000 m




✧ *The best visual depiction of the combined effect of the biological pump & ocean circulation I have seen!*


From Key et al. (2005) in Emerson & Hedges (2007), Fig. 6.16.

*** Ended Here - 2/10/09 ***

Hard vs. Soft Parts







Coccolithophorid (CaCO_3)



Diatom (SiO_2)

- Some organisms precipitate inorganic shells ("hard parts") out of calcium carbonate & silica, the overwhelming majority being coccolithophorids (CaCO_3) & diatoms (SiO_2)
- On average **22 moles** of C as CaCO_3 are precipitated for every **106 moles** of C converted into **phytoplankton biomass** ("soft parts")
- Results in a Redfield Ratio of $\text{C}_{\text{org}}:\text{N}:\text{P}:\text{C}_{\text{CaCO}_3}:\text{O}_2 = 106:16:1:22:-138$
- Production of CaCO_3 is important in air-sea CO_2 exchange because it removes carbon *and* **alkalinity** from surface water & transfers it to deep sea

Globally the Production of CaCO_3 is 0.6 Gt C/yr (~3% of global marine C fixation)



- SeaWiFS images & Composite
- 50% reduction in coccolith blooms expected in N. Atl (their largest habitat) by 2100 A.D. owing to ocean acidification


Iglesias-Rodriguez et al. (2002) *Glob. Biogeochem. Cycl.* Vol. 16(4): 1100, doi: 10.1029/GB001454

Coccolithophorids & Air-Sea CO₂ Exchange (c.)

- Coccolithophorids alter the $C_{\text{organic}}:C_{\text{carbonate}}$ or "rain ratio" & increase surface pCO₂ during calcification, producing one molecule of CO₂ for each molecule of CaCO₃ fixed.
- Globally, the rain ratio from the surface ocean is ~ 4:1
- This ratio is largely controlled by the dominant taxon fixing carbon such that a shift in the phytoplankton community structure from calcifiers to silicifiers would affect the capacity of the biological pump
- For example, during a bloom of coccolithophorids, photosynthesis: calcification & the rain ratio can approach 1:1, & the effect of such high calcification rates has been found to change the air-sea gradient of CO₂.
- Considering that coccolithophorid blooms are responsible for up to 80% of surface ocean calcification, the 50% predicted decrease in potential surface coccolithophorid bloom areal extent may potentially lead to a significant increase in the POC:PIC ratio.
- By 2050 an increase in dissolved CO₂ & a decrease in the concentration of CO₃²⁻ will result in an increase in CaCO₃ dissolution.
- A decrease in calcification is a short-term negative feedback in the global carbon cycle.


IGLESIAS-RODRIGUEZ ET AL. (2002): COCCOLITHOPHORIDS IN OCEAN CARBON CYCLE MODELS

Ratio of Hard to Soft Parts Produced & Exported from the Surface Ocean Influences pCO₂




- Vector diagrams demonstrate the effect of biological & physical processes on surface ocean pCO₂:

- Gas exchange
 - Proportionally changes ocean DIC w/ no effect on alkalinity
- "Soft tissue pump"
 - Photosynthesis reduces DIC & increases alkalinity with a slope of C/N=-106/16=-6.6*
- "Carbonate pump"
 - CaCO₃ production reduces alkalinity by 2 units per unit decrease in DIC b/c CO₃²⁻ contributes 2 moles of alkalinity per mole of DIC




*Debate continues over the "true" Redfield ratio, with this book taking C:N = 117:16

Sarmiento & Gruber (2007) Fig. 8.3.5



Global Ocean 1° Productivity



- Why is productivity so low in subtropical gyres?
- Why is it so high in high latitude oceans? Along equator? On coastal margins?

Ocean color/ SeaWifs: http://oceancolor.gsfc.nasa.gov/SeaWifs/TEACHERS/sanctuary_7.html