



The diagrams at right illustrate the refraction of a wave as it travels from one medium (Region I) to another (Region II.) The wavelength of the refracted waves (λ_r) is twice the wavelength of the incident waves (λ_i).

- A. Determine the ratio of the wave speed in region I to the wave speed in region II. Explain your reasoning.




- B. In the diagram at right, sketch rays indicating the direction in which the waves are propagating. Indicate the angle of incidence (θ_i) and the angle of refraction (θ_r). (Recall, these angles are measured between the direction of propagation and the normal to the boundary.)



- C. The distance d shown in the figure above is the distance between two adjacent crests of the incident (and refracted) waves on the boundary between the two media. As the angle θ_i increases, does d increase, decrease, or remain the same? Explain your reasoning. (You may want to include a sketch to support your answer.)
- D. As the angle θ_i increases does the angle θ_r increase, decrease, or remain the same? Base your answer on your response to part C.
- E. What is the minimum possible value of d for which the crests of the incident waves are aligned with those of the refracted waves? Explain how you can tell. (*Hint:* What is the direction of propagation for the refracted waves when d has its minimum value?)

- F. Sketch the wavefronts for the situation in which the distance \mathbf{d} has its minimum value. Include rays indicating the direction of propagation for the waves in your sketch.



- G. Calculate the angle θ_i for which \mathbf{d} has its minimum value. Show all work.
- H. Is it possible to have a transmitted wave for angles of incidence greater than the angle you have found in G? Explain why or why not.